



Formulary of geodesics of the projected averaged Kepler Hamiltonian

Jean-Baptiste Caillau

► To cite this version:

Jean-Baptiste Caillau. Formulary of geodesics of the projected averaged Kepler Hamiltonian. 2007.
hal-00130214v3

HAL Id: hal-00130214

<https://hal.science/hal-00130214v3>

Preprint submitted on 15 Feb 2007

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Ecole Nationale Supérieure
d'Electrotechnique, Electronique
Informatique, Hydraulique et Télécom
Institut de Recherche en Informatique de Toulouse

**Formulary of geodesics of the projected
averaged Kepler Hamiltonian**

J.-B. Caillau

Équipe Algorithmes Parallèles et Optimisation
ENSEEIHT–IRIT (UMR CNRS 5505)
2 rue Camichel, F-31071 Toulouse

www.n7.fr/apo

Rapport Technique **RT/APO/07/2**

Formulary of geodesics of the projected averaged Kepler Hamiltonian*

J.-B. Caillau[†]

February 2007

Abstract

This short note gives the quadratures of the geodesics of the averaged Hamiltonian of the controlled Kepler equation (energy criterion) projected on \mathbf{S}^2 . The endpoints of the corresponding cut locus are also deduced, as well as the injectivity radius of the associated Riemannian metric on the 2-sphere.

Keywords. Averaged Kepler equation, geodesics on \mathbf{S}^2 , cut locus.

Classification AMS. 49K15, 70Q05

The Hamiltonian is

$$H = \frac{1}{2} \left(\frac{p_\theta^2}{G(\varphi)} + p_\varphi^2 \right)$$

with

$$G(\varphi) = \frac{\sin^2 \varphi}{1 - (1 - \mu^2) \sin^2 \varphi}.$$

Herebefore, μ is a parameter which is equal to $1/\sqrt{5}$ in the case of the Kepler equation [1, 2, 3, 4]. The system is clearly integrable, and the geodesics on the level set $\{H = 1/2\}$ of the corresponding Riemannian metric are as parameterized by p_θ (note that θ is cyclic) as follows:

$$\varphi = \operatorname{asin} \sqrt{\frac{(1+b) - (1-b) \cos(a(t-t_1))}{2}},$$

and

$$\theta = \operatorname{sign}(p_\theta) \left[\operatorname{atan} \left(\frac{\tan(a(t-t_1)/2)}{\sqrt{b}} \right) \right]_0^t - (1 - \mu^2) p_\theta t.$$

These two quadratures are valid for t in $[t_1, t_1 + T/4]$ with nonnegative p_{φ_0} , and extended on $[t_1, t_1 + T]$ by using the obvious symmetries of the Hamiltonian (a $-2t_1$ time translation gives the case when p_{φ_0} is negative). The coordinate

*Travail effectué dans le cadre du Réseau d'Excellence HYCON, contrat numéro FP6-IST-511368.

[†]ENSEEIH-IRIT (UMR CNRS 5505), 2 rue Camichel, F-31071 Toulouse (caillau@n7.fr).

φ and the time derivative $\dot{\theta}$ are T -periodic and hence clearly extended to the whole real line. The period T , the time t_1 , and the parameters a and b have the following expression (θ_0 is assumed to be zero by symmetry):

$$\begin{aligned} a &= 2\sqrt{1 + (1 - \mu^2)p_\theta^2}, \\ b &= p_\theta^2 / (1 + (1 - \mu^2)p_\theta^2), \\ t_1 &= (1/a) \left[-\pi/2 - \operatorname{asin} \left(\frac{2\sin^2 \varphi_0 - (1 + b)}{1 - b} \right) \right], \\ T &= 4\pi/a. \end{aligned}$$

The cut locus [2] of the Riemannian metric is then easily deduced. For symmetry reasons, it is included in the antipodal parallel $\{\pi - \varphi_0\}$, and the coordinates of its left endpoint are

$$\begin{aligned} \theta_l &= \pi(1 - (1 - \mu^2)\sin \varphi_0), \\ \varphi_l &= \pi - \varphi_0. \end{aligned}$$

The time, that is the distance to the cut locus is

$$t_l = \pi \sqrt{1 - (1 - \mu^2)\sin^2 \varphi_0},$$

which gives the injectivity radius of the Riemannian metric $G(\varphi)d\theta^2 + d\varphi^2$ on \mathbf{S}^2 , that is the infimum of distances from a point to its cut locus (clearly reached on the equator, for $\varphi_0 = \pi/2$),

$$i(\mathbf{S}^2) = \mu\pi.$$

As consequence of the computations, any geodesic is—up to a rotation in θ —a *pseudo-equator*, that is a geodesic generated by $p_\varphi = 0$ on $\{H = 1/2\}$. In Kepler case, every pseudo-equator starting from a rational initial eccentricity is closed. There exist five simple closed geodesic modulo rotations with respect to θ , and the shortest closed geodesics are the meridians whose length is 2π .

References

- [1] B. Bonnard and J.-B. Caillau. A global optimality result with application to orbital transfer. *C. R. Acad. Sci. Paris, Ser. I*, submitted.
- [2] B. Bonnard and J.-B. Caillau. Conjugate and cut loci in averaged orbital transfer. *C. R. Acad. Sci. Paris, Ser. I*, submitted.
- [3] B. Bonnard and J.-B. Caillau. Riemannian metric of the averaged energy minimization problem in orbital transfer with low thrust. *An. I. H. Poincaré*, to appear.
- [4] J.-B. Caillau. Sur la géométrie des transferts orbitaux. Habilitation, Insitut National Polytechnique de Toulouse, 2006.